

Universität des Saarlandes



Fachrichtung 6.1 – Mathematik

Preprint Nr. 188

**A Regularity Theory for Scalar Local Minimizers of
Splitting-Type Variational Integrals**

Michael Bildhauer, Martin Fuchs and Xiao Zhong

Saarbrücken 2007

A Regularity Theory for Scalar Local Minimizers of Splitting-Type Variational Integrals

Michael Bildhauer

Saarland University
Department of Mathematics
P.O. Box 15 11 50
D-66041 Saarbrücken
Germany
`bibi@math.uni-sb.de`

Martin Fuchs

Saarland University
Department of Mathematics
P.O. Box 15 11 50
D-66041 Saarbrücken
Germany
`fuchs@math.uni-sb.de`

Xiao Zhong

University of Jyväskylä
Department of Mathematics and Statistics
P.O. Box 35
Fin-40014 University of Jyväskylä
Finland
`zhong@maths.jyu.fi`

Edited by
FR 6.1 – Mathematik
Universität des Saarlandes
Postfach 15 11 50
66041 Saarbrücken
Germany

Fax: + 49 681 302 4443
e-Mail: preprint@math.uni-sb.de
WWW: <http://www.math.uni-sb.de/>

Abstract

Starting from Giaquinta's counterexample [Gi] we introduce the class of splitting functionals being of (p, q) -growth with exponents $p \leq q < \infty$ and show for the scalar case that locally bounded local minimizers are of class $C^{1,\mu}$. Note that to our knowledge the only $C^{1,\mu}$ -results without imposing a relation between p and q concern the case of two independent variables as it is outlined in Marcellini's paper [Ma1], Theorem A, and later on in the work of Fusco and Sbordone [FS], Theorem 4.2.

1 Introduction

In 1987 Giaquinta [Gi] showed that the function

$$u_0(x) := \sqrt{\frac{n-4}{24}} x_n / \sqrt{x_1^2 + \dots + x_{n-1}^2},$$

$$x \in B_1(0) = \{y \in \mathbb{R}^n : |y| < 1\},$$

is a local minimizer of the energy

$$J[w] = \int_{B_1(0)} \left[\sum_{i=1}^{n-1} (\partial_i w)^2 + \frac{1}{2} (\partial_n w)^4 \right] dx, \quad (1.1)$$

provided that $n \geq 6$. Since u_0 is unbounded, Giaquinta's example clearly demonstrates that for anisotropic variational integrals in general no regularity results for local minimizers can be expected, and this even concerns the scalar situation! So one may ask for an admissible range of anisotropy implying that local minimizers are locally bounded (and even share a higher degree of regularity) or one may try to compensate the anisotropic structure of the integrand by adding certain natural hypothesis on the behaviour of the local minimizer leading to its regularity. But let us first have a closer look at Giaquinta's energy J . If we write $z = (\tilde{z}, z_n)$, $\tilde{z} := (z_1, \dots, z_{n-1})$, for elements z of \mathbb{R}^n , $n \geq 2$, then the energy density occurring in (1.1) is of splitting form, i.e. we have an integrand $F : \mathbb{R}^n \rightarrow [0, \infty)$ such that

$$F(z) = f(\tilde{z}) + g(z_n) \quad (1.2)$$

with C^2 -functions $f : \mathbb{R}^{n-1} \rightarrow [0, \infty)$, $g : \mathbb{R} \rightarrow [0, \infty)$ satisfying with exponents $1 < p \leq q < \infty$ and with constants $\lambda, \Lambda > 0$ the ellipticity conditions ($y, z \in \mathbb{R}^n$)

$$\lambda(1 + |\tilde{z}|^2)^{\frac{p-2}{2}} |\tilde{y}|^2 \leq D^2 f(\tilde{z})(\tilde{y}, \tilde{y}) \leq \Lambda(1 + |\tilde{z}|^2)^{\frac{p-2}{2}} |\tilde{y}|^2, \quad (1.3)$$

$$\lambda(1 + |z_n|^2)^{\frac{q-2}{2}} |y_n|^2 \leq D^2 g(z_n)(y_n, y_n) \leq \Lambda(1 + |z_n|^2)^{\frac{q-2}{2}} |y_n|^2. \quad (1.4)$$

Of course (1.4) could be stated in a simpler form but (1.2) just serves as a model case: in fact we could consider any decomposition $z = (z^{(1)}, z^{(2)})$ of the vector z and replace (1.2) by $F(z) = f(z^{(1)}) + g(z^{(2)})$ with f and g satisfying the appropriate versions of (1.3) and (1.4). We

AMS Subject Classification: 49N60

Keywords: anisotropic problems, splitting integrals, regularity of minimizers, scalar case

Acknowledgement: A part of this paper was finished during the stay of M. Bildhauer and M. Fuchs at the University of Jyväskylä in November 2006. We like to thank the members of the Dept. of Math. for their hospitality.

also like to remark that for $p \geq 2$ the degenerate variants of (1.3), (1.4) can be considered so that Giaquinta's energy (1.1) is included. Now, if Ω is an open set in \mathbb{R}^n , we let

$$I[w, \Omega] = \int_{\Omega} F(\nabla w) \, dx, \quad (1.5)$$

and since (1.3), (1.4) imply the growth estimate

$$a|z|^p - b \leq F(z) \leq A|z|^q + B, \quad z \in \mathbb{R}^n, \quad (1.6)$$

with constants $a, A > 0, b, B \geq 0$, it is natural to call a function u from the local Sobolev space $W_{p,loc}^1(\Omega)$ (see [Ad] for a definition) a local minimizer of I from (1.5) iff $I[u, \Omega'] < \infty$ and $I[u, \Omega'] \leq I[w, \Omega']$ for any open set Ω' s.t. $\Omega' \Subset \Omega$ and for all $w \in W_{p,loc}^1(\Omega)$ such that $\text{spt}(u - w) \Subset \Omega'$.

Coming back to the regularity problem for local minimizers u , it turns out that conditions of the form

$$q \leq c(n)p \quad (1.7)$$

are sufficient for the local boundedness of the function u and also for its higher regularity. Here $c(n)$ is a constant depending on n giving rather large values if n is small but with the unpleasant property $c(n) \rightarrow 1$ as $n \rightarrow \infty$. We mention the contributions of Fusco and Sbordone [FS], Marcellini [Ma1], [Ma2] and Hong [Ho], where one also finds further references. It should be remarked that results of this type usually do not refer to a splitting structure of the integrand F as stated in (1.2): in place of this one works with (1.6) combined with an appropriate ellipticity condition, or one just requires that

$$\lambda(1 + |z|^2)^{\frac{p-2}{2}}|y|^2 \leq D^2F(z)(y, y) \leq \Lambda(1 + |z|^2)^{\frac{q-2}{2}}|y|^2, \quad (1.8)$$

which obviously implies the validity of (1.6). The reader should note that for integrands F satisfying (1.2) the conditions (1.3), (1.4) do not imply (1.8), and that conversely (1.3), (1.4) can not be deduced from (1.8).

Let us now look at a special situation for which it is possible to improve (1.7): if Ω is a bounded domain and if u is an $I[\cdot, \Omega]$ -minimizer for boundary data $\bar{u} \in L^\infty(\Omega)$, which means that we are given $\bar{u} \in W_p^1(\Omega) \cap L^\infty(\Omega)$ such that $I[\bar{u}, \Omega] < \infty$ and where the boundary condition has to be understood in the sense that $u - \bar{u} \in \overset{\circ}{W}_p^1(\Omega)$, then the maximum-principle implies $u \in L^\infty(\Omega)$. From this point of view it makes sense to study local minimizers from the space $L_{loc}^\infty(\Omega)$. Assuming this together with (1.8), Choe [Ch] proved the smoothness of u under the dimensionless condition $q < p + 1$, which was replaced by $q < p + 2$ in [Bi] (see also [ELM]) and [BF1]. If the integrand F is of splitting type satisfying (1.3), (1.4) with $q \leq 2p$, then in the recent paper [BFZ] we could show the differentiability of local minimizers $u \in L_{loc}^\infty(\Omega)$ which leads to a partial improvement of earlier results of Ural'tseva and Urdaletova [UU]. In the present note we are going to apply new methods leading to the following results:

Theorem 1.1. *Let $1 < p \leq q < \infty$, and let F satisfy (1.2), (1.3) and (1.4). Consider a local $I[\cdot, \Omega]$ -minimizer u of class $W_{p,loc}^1(\Omega) \cap L_{loc}^\infty(\Omega)$. Then we have:*

- i) $\nabla u \in L_{loc}^m(\Omega; \mathbb{R}^n)$ for any finite exponent m .
- ii) If in addition $p \geq 2$, then $u \in C^{1,\mu}(\Omega)$ holds for all $\mu < 1$.

Remark 1.1. *If the case of degenerate ellipticity is considered in (1.3) and (1.4), then we have Theorem 1.1, i) under the restriction that $p \geq 2$. In this case we get in Theorem 1.1, ii) that $u \in C^{1,\mu}(\Omega)$ for some $\mu < 1$, the necessary adjustments can be found, for instance, in [BFM].*

Remark 1.2. We like to emphasize again that our results are not limited to the specific decomposition (1.2). With minor modifications we can also discuss the integrand

$$F(z) = \sum_{i=1}^n (1 + z_i^2)^{p_i/2}, \quad z \in \mathbb{R}^n,$$

with exponents $1 < p_i < \infty$. Alternatively we may consider a decomposition

$$F(z) := F^{(1)}(z^{(1)}) + F^{(2)}(z^{(2)}),$$

where for example $z^{(1)} := (z_1, \dots, z_k)$, $z^{(2)} := (z_{k+1}, \dots, z_n)$ with $1 \leq k < n$ and where $F^{(1)}$ and $F^{(2)}$ satisfy ellipticity conditions like (1.3) and (1.4) with exponents p_1 and p_2 .

Remark 1.3. Going through the proof of Theorem 1.1 one easily checks that the explicit additive structure of F itself formulated in (1.2) is not really needed. In fact, if we drop (1.2) and replace (1.3) and (1.4) by the assumption

$$\begin{aligned} \lambda \left[(1 + |\tilde{z}|^2)^{\frac{p-2}{2}} |\tilde{y}|^2 + (1 + |z_n|^2)^{\frac{q-2}{2}} |y_n|^2 \right] &\leq D^2 F(z)(y, y) \\ &\leq \Lambda \left[(1 + |\tilde{z}|^2)^{\frac{p-2}{2}} |\tilde{y}|^2 + (1 + |z_n|^2)^{\frac{q-2}{2}} |y_n|^2 \right], \end{aligned}$$

then we still have our results stated above.

Remark 1.4. It is easy to extend Theorem 1.1 to non-autonomous energies $F(x, z)$, $x \in \overline{\Omega}$, $z \in \mathbb{R}^n$, provided $D_x D_z F(x, z)$ satisfies a natural growth condition.

Remark 1.5. In the case of vector-valued functions we have much weaker results which are summarized in [BF2].

As an application of Theorem 1.1 we consider the following variant of Giaquinta's energy which was introduced by Fusco and Sbordone (see [FS], formula (3.4)): for $q > 2$ and a positive constant c let

$$J_q[w] := \int_{B_1(0)} \left[\sum_{i=1}^{n-1} (\partial_i w)^2 + \frac{2c}{q} |\partial_n w|^q \right] dx.$$

If $n \leq 3$, then according to Theorem 3.1 in [FS] any local J_q -minimizer is locally bounded independent of the value of q , whereas Fusco and Sbordone obtain $C^{1,\alpha}$ -regularity for the non-degenerate variant of J_q if $q \leq 2n/(n-2)$ is satisfied (see Theorem 4.2 in [FS]), i.e. they require the bound $q \leq 6$ in the 3D-case. But Theorem 1.1 combined with the local boundedness result of [FS] shows

Corollary 1.1. *If $n \leq 3$ and if $q \geq 2$ is arbitrary, then all local minima of the functional J_q belong to the class $C^{1,\alpha}(B_1(0))$ for an exponent $\alpha > 0$.*

We leave it to the reader to give a version of this corollary valid for more general functionals on three-dimensional domains.

2 Proof of Theorem 1.1

Obviously Theorem 1.1, ii) follows from part i) combined with Lemma 3.1. To prove i) we fix a ball $B := B_R(x_0)$ with compact closure in Ω and consider the mollification $(u)_\epsilon$ with a small radius $\epsilon > 0$. Let u_ϵ denote the unique Lipschitz function minimizing $I[\cdot, B]$ in the class of all Lipschitz mapping $\overline{B} \rightarrow \mathbb{R}$ for boundary values $(u)_\epsilon$, i.e. u_ϵ denotes the Hilbert-Haar solution (see, e.g., [MM], Theorem 4, p. 162). The main properties of this approximation are summarized in

Lemma 2.1. i) *Passing to the limit $\epsilon \rightarrow 0$ we have*

$$u_\epsilon \rightharpoonup u \quad \text{in } W_p^1(B), \quad \int_B F(\nabla u_\epsilon) \, dx \rightarrow \int_B F(\nabla u) \, dx.$$

ii) $\|u_\epsilon\|_{L^\infty(B)}$ *is bounded independent of ϵ .*

Proof. i) The minimality of u_ϵ implies

$$\int_B F(\nabla u_\epsilon) \, dx \leq \int_B F(\nabla(u)_\epsilon) \, dx,$$

and from Jensen's inequality we deduce

$$\int_B F(\nabla(u)_\epsilon) \, dx \leq \int_B F(\nabla u) \, dx + O(\epsilon)$$

with $O(\epsilon) \rightarrow 0$ as $\epsilon \rightarrow 0$. Due to the growth of F we find $\sup_{\epsilon>0} \|u_\epsilon\|_{W_p^1(B)} < \infty$, so that $u_\epsilon \rightharpoonup \bar{u}$ in $W_p^1(B)$ for some function \bar{u} from this class. By lower semicontinuity it holds

$$\int_B F(\nabla \bar{u}) \, dx \leq \liminf_{\epsilon \rightarrow 0} \int_B F(\nabla u_\epsilon) \, dx,$$

hence

$$\int_B F(\nabla \bar{u}) \, dx \leq \int_B F(\nabla u) \, dx.$$

On the other hand we have $u_\epsilon - (u)_\epsilon \in \mathring{W}_p^1(B)$, which means $\bar{u} - u \in \mathring{W}_p^1(B)$. Clearly u minimizes $I[\cdot, B]$ w.r.t. its own boundary values, and the strict convexity of F implies $\bar{u} = u$.

ii) Let $v := \min\{u_\epsilon, \sup_{\partial B}(u)_\epsilon\}$. Then v is Lipschitz on B with trace $(u)_\epsilon$, thus

$$I[u_\epsilon, B] \leq I[v, B]$$

and in conclusion $u_\epsilon \leq \sup_{\partial B}(u)_\epsilon$. In the same way we obtain $u_\epsilon \geq \min_{\partial B}(u)_\epsilon$, and since $u \in L^\infty(B)$, the claim follows from these estimates. \square

Lemma 2.2. *The functions u_ϵ are of class $C^{1,\mu}(B) \cap W_{2,loc}^2(B)$ for any $\mu < 1$.*

Proof. This result is standard and can be found in the textbook of Massari and Miranda, [MM], Theorem 5, p. 166. The first lines of the proof of Lemma 3.1 summarize the idea observing that by definition of u_ϵ we here already have $\nabla u_\epsilon \in L^\infty(B; \mathbb{R}^n)$. \square

Lemma 2.3. *(Variants of Caccioppoli's inequality) For any numbers $\alpha, \beta \geq 0$ and for all $\eta \in C_0^\infty(B)$ s.t. $0 \leq \eta \leq 1$ we have*

$$\int_B D^2 F(\nabla u_\epsilon) (\partial_n \nabla u_\epsilon, \partial_n \nabla u_\epsilon) \Gamma_{n,\epsilon}^{\frac{\alpha}{2}} \eta^2 \, dx \leq c(\alpha) \int_\Omega D^2 F(\nabla u_\epsilon) (\nabla \eta, \nabla \eta) \Gamma_{n,\epsilon}^{\frac{\alpha+2}{2}} \, dx, \quad (2.1)$$

$$\int_B D^2 F(\nabla u_\epsilon) (\partial_\gamma \nabla u_\epsilon, \partial_\gamma \nabla u_\epsilon) \tilde{\Gamma}_\epsilon^{\frac{\beta}{2}} \eta^2 \, dx \leq c(\beta) \int_B D^2 F(\nabla u_\epsilon) (\nabla \eta, \nabla \eta) \tilde{\Gamma}_\epsilon^{\frac{\beta+2}{2}} \, dx. \quad (2.2)$$

In (2.2) (and in what follows) we always take the sum w.r.t. γ from 1 to $n-1$. $c(\alpha)$, $c(\beta)$ denote positive constants independent of ϵ , and we have set: $\Gamma_{n,\epsilon} = 1 + (\partial_n u_\epsilon)^2$, $\tilde{\Gamma}_\epsilon = 1 + |\tilde{\nabla} u_\epsilon|^2$, $\tilde{\nabla} := (\partial_1, \dots, \partial_{n-1})$.

Proof. See, e.g., the beginnings of Section 3 and Section 4 of [BFZ]. We like to remark that by Lemma 2.2 any first partial derivative of u_ϵ is a solution of an uniformly elliptic equation with continuous coefficients so that by standard potential theory we get $\partial_k u_\epsilon \in W_{m,loc}^1(B)$ for any $m < \infty$ and all $k = 1, \dots, n$, i.e. $u_\epsilon \in W_{m,loc}^2(B)$ for all $m < \infty$. For this reason the calculations leading to (2.1) and (2.2), which were carried out in [BFZ] for a different type of approximation, can be justified in the present setting. \square

Now, with $\alpha \geq 0$ and $\eta \in C_0^\infty(B)$, $0 \leq \eta \leq 1$, being fixed for the moment we consider the expression

$$\int_B \eta^2 \Gamma_{n,\epsilon}^{\frac{q+2+\alpha}{2}} dx = \int_B \eta^2 \Gamma_{n,\epsilon}^{\frac{q+\alpha}{2}} dx + \int_B \partial_n u_\epsilon \partial_n u_\epsilon \eta^2 \Gamma_{n,\epsilon}^{\frac{q+\alpha}{2}} dx \quad (2.3)$$

which is well defined by the Lipschitz continuity of u_ϵ , and according to Lemma 2.2 we are allowed to integrate by parts in the second term on the r.h.s. of (2.3), i.e.

$$\int_B \partial_n u_\epsilon \partial_n u_\epsilon \eta^2 \Gamma_{n,\epsilon}^{\frac{q+\alpha}{2}} dx = - \int_B u_\epsilon \partial_n \left[\partial_n u_\epsilon \eta^2 \Gamma_{n,\epsilon}^{\frac{q+\alpha}{2}} \right] dx.$$

Using Lemma 2.1 ii), we see that

$$\begin{aligned} \int_B \eta^2 \Gamma_{n,\epsilon}^{\frac{q+\alpha+2}{2}} dx &\leq c \left[\int_B \eta^2 \Gamma_{n,\epsilon}^{\frac{q+\alpha}{2}} dx + \int_B \eta |\nabla \eta| |\partial_n u_\epsilon| \Gamma_{n,\epsilon}^{\frac{q+\alpha}{2}} dx + \int_B \eta^2 |\partial_n \partial_n u_\epsilon| \Gamma_{n,\epsilon}^{\frac{q+\alpha}{2}} dx \right] \\ &=: c \left[\int_B \eta^2 \Gamma_{n,\epsilon}^{\frac{q+\alpha}{2}} + I_1 + I_2 \right] \end{aligned} \quad (2.4)$$

with a constant c depending also on α but being independent of ϵ and η . In order to handle I_1 and I_2 we will apply Young's inequality with small parameter τ which enables us to absorb terms in the l.h.s. of (2.4). We have

$$\begin{aligned} I_1 &\leq \int_B \eta |\nabla \eta| \Gamma_{n,\epsilon}^{\frac{q+1+\alpha}{2}} dx \leq \tau \int_B \eta^2 \Gamma_{n,\epsilon}^{\frac{q+2+\alpha}{2}} dx + c(\tau) \int_B |\nabla \eta|^2 \Gamma_{n,\epsilon}^{\frac{q+\alpha}{2}} dx, \\ I_2 &= \int_B \eta^2 |\partial_n \partial_n u_\epsilon| \Gamma_{n,\epsilon}^{\frac{q-2+\alpha}{4}} \Gamma_{n,\epsilon}^{\frac{q+2+\alpha}{4}} dx \\ &\leq \tau \int_B \Gamma_{n,\epsilon}^{\frac{q+2+\alpha}{2}} \eta^2 dx + c(\tau) \int_B \eta^2 |\partial_n \partial_n u_\epsilon|^2 \Gamma_{n,\epsilon}^{\frac{q-2+\alpha}{2}} dx, \end{aligned}$$

which implies by (2.4)

$$\begin{aligned} \int_B \eta^2 \Gamma_{n,\epsilon}^{\frac{q+2+\alpha}{2}} dx &\leq c \left[\int_B (\eta^2 + |\nabla \eta|^2) \Gamma_{n,\epsilon}^{\frac{q+\alpha}{2}} dx + J \right], \\ J &:= \int_B \eta^2 |\partial_n \partial_n u_\epsilon|^2 \Gamma_{n,\epsilon}^{\frac{q-2+\alpha}{2}} dx. \end{aligned} \quad (2.5)$$

Then we use (1.3), (1.4) and (2.1) to obtain

$$\begin{aligned} J &\leq c \int_B \eta^2 D^2 F(\nabla u_\epsilon) (\partial_n \nabla u_\epsilon, \partial_n \nabla u_\epsilon) \Gamma_{n,\epsilon}^{\frac{\alpha}{2}} dx \\ &\leq c \int_B D^2 F(\nabla u_\epsilon) (\nabla \eta, \nabla \eta) \Gamma_{n,\epsilon}^{\frac{\alpha+2}{2}} dx \\ &\leq c \left[\int_B \tilde{\Gamma}_\epsilon^{\frac{p-2}{2}} \Gamma_{n,\epsilon}^{\frac{\alpha+2}{2}} |\nabla \eta|^2 dx + \int_B |\nabla \eta|^2 \Gamma_{n,\epsilon}^{\frac{q+\alpha}{2}} dx \right]. \end{aligned}$$

Inserting this into (2.5) it follows

$$\int_B \eta^2 \Gamma_{n,\epsilon}^{\frac{q+2+\alpha}{2}} dx \leq c \left[\int_B (\eta^2 + |\nabla \eta|^2) \Gamma_{n,\epsilon}^{\frac{q+\alpha}{2}} dx + \int_B \tilde{\Gamma}_\epsilon^{\frac{p-2}{2}} \Gamma_{n,\epsilon}^{\frac{\alpha+2}{2}} |\nabla \eta|^2 dx \right]. \quad (2.6)$$

Next we let $\beta \geq 0$ and fix η as above. Then (recall that the sum is taken w.r.t. $\gamma = 1, \dots, n-1$) we consider the expression

$$\int_B \eta^2 \tilde{\Gamma}_\epsilon^{\frac{p+2+\beta}{2}} dx,$$

perform an integration by parts in

$$\int_B \partial_\gamma u_\epsilon \partial_\gamma u_\epsilon \eta^2 \tilde{\Gamma}_\epsilon^{\frac{p+\beta}{2}} dx$$

and carry out analogous calculations leading to the following variant of (2.5):

$$\int_B \eta^2 \tilde{\Gamma}_\epsilon^{\frac{p+2+\beta}{2}} dx \leq c \left[\int_B (\eta^2 + |\nabla \eta|^2) \tilde{\Gamma}_\epsilon^{\frac{p+\beta}{2}} dx + \int_B \eta^2 |\tilde{\nabla}^2 u_\epsilon|^2 \tilde{\Gamma}_\epsilon^{\frac{p-2+\beta}{2}} dx \right]. \quad (2.7)$$

The second item on the r.h.s. of (2.7) is handled in the same manner as J from above, i.e. this expression is bounded by

$$c \int_B \eta^2 D^2 F(\nabla u_\epsilon) (\partial_\gamma \nabla u_\epsilon, \partial_\gamma \nabla u_\epsilon) \tilde{\Gamma}_\epsilon^{\frac{\beta}{2}} dx \leq c \left[\int_B |\nabla \eta|^2 \tilde{\Gamma}_\epsilon^{\frac{p+\beta}{2}} dx + \int_B |\nabla \eta|^2 \Gamma_{n,\epsilon}^{\frac{q-2}{2}} \tilde{\Gamma}_\epsilon^{\frac{\beta+2}{2}} dx \right],$$

thus we get

$$\int_B \eta^2 \tilde{\Gamma}_\epsilon^{\frac{p+2+\beta}{2}} dx \leq c \left[\int_B (\eta^2 + |\nabla \eta|^2) \tilde{\Gamma}_\epsilon^{\frac{p+\beta}{2}} + \int_B |\nabla \eta|^2 \Gamma_{n,\epsilon}^{\frac{q-2}{2}} \tilde{\Gamma}_\epsilon^{\frac{\beta+2}{2}} dx \right]. \quad (2.8)$$

We return to (2.6) replacing η by η^k for a large number $k \in \mathbb{N}$, apply Young's inequality on the r.h.s. in order to get terms of the form $\tau \int_B \eta^{2k} \Gamma_{n,\epsilon}^{\frac{q+2+\alpha}{2}} dx$ which can be absorbed in the l.h.s. and get

$$\begin{aligned} \int_B \eta^{2k} \Gamma_{n,\epsilon}^{\frac{q+2+\alpha}{2}} dx &\leq c \left[\int_B \left(\eta^{2k} + |\nabla \eta|^{q+2+\alpha} \eta^{2k-(q+2+\alpha)} \right) dx \right. \\ &\quad \left. + \int_B |\nabla \eta|^{\frac{2}{q}(\alpha+q+2)} \eta^{2k-\frac{2}{q}(\alpha+q+2)} \tilde{\Gamma}_\epsilon^{\frac{p-2}{2} \frac{\alpha+q+2}{q}} dx \right]. \end{aligned} \quad (2.9)$$

In the same way (2.8) implies

$$\begin{aligned} \int_B \eta^{2k} \tilde{\Gamma}_\epsilon^{\frac{p+2+\beta}{2}} dx &\leq c \left[\int_B \left(\eta^{2k} + |\nabla \eta|^{p+2+\beta} \eta^{2k-(p+2+\beta)} \right) dx \right. \\ &\quad \left. + \int_B |\nabla \eta|^{\frac{2}{p}(\beta+2+p)} \eta^{2k-\frac{2}{p}(\beta+2+p)} \Gamma_{n,\epsilon}^{\frac{q-2}{2} \frac{p+2+\beta}{p}} dx \right]. \end{aligned} \quad (2.10)$$

Let us introduce the quantities

$$\begin{aligned}
a(\alpha, \rho) &:= \int_{B_\rho} \Gamma_{n,\epsilon}^{\frac{q+2+\alpha}{2}} dx, \\
A(\alpha, \rho) &:= \int_{B_\rho} \tilde{\Gamma}_\epsilon^{\frac{p-2}{2} \frac{\alpha+q+2}{q}} dx, \\
b(\beta, \rho) &:= \int_{B_\rho} \tilde{\Gamma}_\epsilon^{\frac{p+2+\beta}{2}}, \\
B(\beta, \rho) &:= \int_{B_\rho} \Gamma_{n,\epsilon}^{\frac{q-2}{2} \frac{p+2+\beta}{p}} dx,
\end{aligned}$$

where $B_\rho := B_\rho(x_0)$ with $\rho < R$. If $p \leq 2$, then (2.9) immediately implies the uniform local integrability of $\partial_n u_\epsilon$ for any finite exponent m , and using this information in (2.10) we get the same result for $\tilde{\nabla} u_\epsilon$. So let us assume $p > 2$. Then we have

$$A(\alpha_0, \rho) \leq c(\rho) < \infty \quad (2.11)$$

for a constant $c(\rho)$ going to infinity as $\rho \uparrow R$ but being independent of ϵ , provided we require

$$\frac{p-2}{2} \frac{q+\alpha_0+2}{q} \leq \frac{p}{2} \quad (2.12)$$

and quote Lemma 2.1 i). Note that on account of $(p-2)(q+2) < pq$ (2.12) holds for suitable positive numbers α_0 and we may choose

$$\alpha_0 := \frac{pq}{p-2} - (q+2) > 0. \quad (2.13)$$

(2.11) combined with (2.9) shows that

$$a(\alpha_0, \rho) \leq c(\rho). \quad (2.14)$$

Now we select β_0 such that

$$\frac{q-2}{2} \frac{p+\beta_0+2}{p} = \frac{q+2+\alpha_0}{2}, \quad (2.15)$$

hence $B(\beta_0, \rho) \leq c(\rho)$ on account of (2.14) and the definition of $B(\beta, \rho)$. Note that by the definition of α_0 we have

$$\frac{q-2}{2} \frac{p+2}{2} < \frac{q+\alpha_0+2}{2}$$

which is equivalent to $-4q - 2p^2 + 8 < 0$ (recall that we assume $p > 2$), thus the solution β_0 of (2.15) is a positive number. Returning to (2.11) we have shown that

$$A(\alpha_l, \rho) + B(\beta_l, \rho) \leq c_l(\rho) \quad (2.16_l)$$

at least for $l = 0$. Suppose that $l \geq 1$ and that (2.16 $_{l-1}$) is valid. We then like to prove (2.16 $_l$) for suitable exponents α_l, β_l . First, $B(\beta_{l-1}, \rho) \leq c_{l-1}(\rho)$ together with (2.10) gives $b(\beta_{l-1}, \rho) \leq c(\rho)$ for a new constant (depending on l) so that

$$A(\alpha_l, \rho) \leq c(\rho), \quad (2.17)$$

provided

$$\frac{p-2}{2} \frac{\alpha_l+q+2}{q} \leq \frac{p+2+\beta_{l-1}}{2},$$

and we may define

$$\begin{aligned}
\alpha_l &= \frac{q}{p-2}(\beta_{l-1} + p + 2) - (q + 2) \\
&= \frac{q}{p-2}\beta_{l-1} + \frac{q(p+2)}{p-2} - (q + 2) \\
&= \frac{q}{p-2}\beta_{l-1} + \frac{4q - 2p + 4}{p-2}
\end{aligned} \tag{2.18}$$

to ensure (2.17). (2.17) together with (2.9) implies $a(\alpha_l, \rho) \leq c(\rho)$ so that

$$B(\beta_l, \rho) \leq c(\rho),$$

if β_l satisfies

$$\frac{q-2}{2} \frac{p+2+\beta_l}{p} = \frac{q+2+\alpha_l}{2},$$

which means

$$\beta_l = \frac{p}{q-2} \alpha_l + \frac{1}{q-2} [4p + 4 - 2q]. \tag{2.19}$$

Inserting (2.18) in (2.19) we see

$$\begin{aligned}
\beta_l &= \frac{pq}{(p-2)(q-2)} \beta_{l-1} + \xi \\
\xi &:= \frac{1}{q-2} [4p + 4 - 2q] + \frac{1}{q-2} \frac{1}{p-2} p [4q - 2p + 4] \\
&= \frac{2p^2 + 2pq + 4q - 8}{(q-2)(p-2)} > 0,
\end{aligned} \tag{2.20}$$

hence the sequence β_l consists of strictly positive numbers. Altogether we have shown the validity of (2.16_l) for any l provided $\{\alpha_l\}$, $\{\beta_l\}$ are defined according to (2.18), (2.20) with the initial values from (2.13) and (2.15). Moreover, since

$$\frac{pq}{(p-2)(q-2)} > 1,$$

we see from (2.20) that $\beta_l \rightarrow \infty$ as $l \rightarrow \infty$, and (2.18) gives the same for α_l which means that

$$\int_{B_\rho} |\nabla u_\epsilon|^m dx \leq c(m, \rho) \tag{2.21}$$

for any $\rho < R$ and all $m < \infty$, the constant being independent of ϵ . Now the claim of Theorem 1.1, i) follows from (2.21) combined with Lemma 2.1, i). \square

3 Some auxiliary results

In this section we will establish a regularity result which might be known but which we could not trace in the literature. As already remarked it will enable us to deduce part ii) of Theorem 1.1 from part i).

Lemma 3.1. *Suppose that $2 \leq s \leq t < \infty$ and suppose that we are given functions $f^*: \mathbb{R}^{n-1} \rightarrow [0, \infty)$, $g^*: \mathbb{R} \rightarrow [0, \infty)$ satisfying (1.3) and (1.4) with exponents s and t respectively. Suppose also that $u^* \in W_{s,loc}^1(\Omega)$ locally minimizes the functional $I^*[\cdot, \Omega]$, where*

$$I^*[w, \Omega] := \int_{\Omega} F^*(\nabla w) dx, \quad F^*(z) := f^*(\tilde{z}) + g^*(z_n).$$

Then, if $\nabla u^ \in L_{loc}^\rho(\Omega, \mathbb{R}^n)$ for any $\rho < \infty$, we have that $u^* \in C^{1,\mu}(\Omega)$ for all $\mu < 1$.*

Proof of Lemma 3.1. It is enough to show that ∇u^* is locally bounded. Then, passing to difference quotients, it is easy to see that $\Delta_h u^*$ is a solution of a uniformly elliptic equation with bounded measurable coefficients, all bounds being independent of h . The De Giorgi-Moser-Nash theory (see [GT]) implies $\Delta_h u \in C^{0,\mu}(\Omega)$ uniformly in h , and the claim follows by passing to the limit $h \rightarrow 0$. For proving the local boundedness of the gradient, we simplify our notation and write u, f, g, F in place of u^*, f^*, g^*, F^* assuming that (1.3) holds for f with exponent $s \geq 2$ and that (1.4) is true for g with exponent $t \geq s$. We first use our assumption

$$\nabla u \in L_{loc}^\rho(\Omega; \mathbb{R}^n) \quad \text{for all } \rho < \infty \quad (3.1)$$

to show that ∇u is weakly differentiable. Let e_γ denote a coordinate direction, $\gamma = 1, \dots, n$, and let

$$\Delta_h^\gamma \Psi(x) := \frac{1}{h} (\Psi(x + h e_\gamma) - \Psi(x)), \quad h \neq 0,$$

denote the corresponding difference quotient of a function Ψ . The minimality of u implies the Euler-equation ($\Omega' \Subset \Omega$)

$$\int_{\Omega'} DF(\nabla u) \cdot \nabla \varphi \, dx = 0 \quad (3.2)$$

valid for $\varphi \in C_0^\infty(\Omega')$ but according to (3.1) φ can be taken from any space $\mathring{W}_\tau^1(\Omega')$, $\tau > 1$, in particular we may choose $\varphi = \Delta_{-h}^\gamma(\eta^2 \Delta_h^\gamma u)$, where $\eta \in C_0^\infty(\Omega)$ is fixed and $|h| \ll 1$ is sufficiently small. From (3.2) we get

$$\int_{\Omega} \Delta_h^\gamma (DF(\nabla u)) \cdot \nabla (\eta^2 \Delta_h^\gamma u) \, dx = 0, \quad (3.3)$$

and if we introduce the bilinear form

$$\mathcal{B}_h^\gamma := \int_0^1 D^2 F(\nabla u + t h \Delta_h^\gamma \nabla u) \, dt,$$

then (3.3) takes the form (no summation w.r.t. γ)

$$\int_{\Omega} \mathcal{B}_h^\gamma (\Delta_h^\gamma \nabla u, \Delta_h^\gamma \nabla u) \eta^2 \, dx = -2 \int_{\Omega} \eta \mathcal{B} (\Delta_h^\gamma \nabla u, \nabla \eta) \Delta_h^\gamma u \, dx. \quad (3.4)$$

Using on the r.h.s. the Cauchy-Schwarz inequality for the bilinear form \mathcal{B} together with Young's inequality we deduce from (3.4) the estimate

$$\int_{\Omega} \mathcal{B}_h^\gamma (\Delta_h^\gamma \nabla u, \Delta_h^\gamma \nabla u) \eta^2 \, dx \leq c \int_{\Omega} \mathcal{B}_h^\gamma (\nabla \eta, \nabla \eta) |\Delta_h^\gamma u|^2 \, dx. \quad (3.5)$$

Elementary properties of the difference quotients in combination with (3.1) show that

$$\int_{\Omega} \mathcal{B}_h^\gamma (\nabla \eta, \nabla \eta) |\Delta_h^\gamma u|^2 \, dx \longrightarrow \int_{\Omega} D^2 F(\nabla u) (\nabla \eta, \nabla \eta) |\partial_\gamma u|^2 \, dx$$

as $h \rightarrow 0$, the limit of course being finite on account of (3.1). Since $s \geq 2$ the l.h.s. of (3.5) is bounded from below by $c \int_{\Omega} \eta^2 |\Delta_h^\gamma \nabla u|^2 \, dx$, hence

$$\int_{\Omega} \eta^2 |\Delta_h^\gamma \nabla u|^2 \, dx \leq c(\eta) < \infty \quad (3.6)$$

for $|h| \ll 1$, and since (3.6) holds for any direction and arbitrary $\eta \in C_0^\infty(\Omega)$, it follows that u is in the space $W_{2,loc}^2(\Omega)$, in particular it holds $\Delta_h^\gamma \nabla u \rightarrow \partial_\gamma \nabla u$ a.e. Since $\mathcal{B}_h^\gamma (\Delta_h^\gamma \nabla u, \Delta_h^\gamma \nabla u) \geq 0$ we may therefore apply Fatou's lemma on the l.h.s. of (3.5) leading to the inequality

$$\int_{\Omega} \eta^2 D^2 F(\nabla u) (\partial_\gamma \nabla u, \partial_\gamma \nabla u) \, dx \leq c \int_{\Omega} D^2 F(\nabla u) (\nabla \eta, \nabla \eta) |\partial_\gamma u|^2 \, dx. \quad (3.7)$$

Note that (3.7) can also be established if $s < 2$, we refer to [Bi], p. 56. Now we fix a ball $B := B_R(x_0)$ with compact closure in Ω . Let $0 < r < \hat{r} < R$ (all balls are centered at x_0), $k > 0$, and define

$$\begin{aligned}\Gamma_n &:= 1 + |\partial_n u|^2, & \Gamma_n^{(h)} &:= 1 + |\Delta_h^n u|^2, \\ \tilde{\Gamma} &:= 1 + |\tilde{\nabla} u|^2, & \tilde{\nabla} u &:= (\partial_1 u, \dots, \partial_{n-1} u).\end{aligned}$$

Then, choosing $\eta \in C_0^\infty(B_{\hat{r}})$, $\varphi := \Delta_{-h}^n(\eta^2 \Delta_h^n u [\Gamma_n^{(h)} - k]^+)$ is admissible in (3.2) leading to

$$\begin{aligned}0 &= \int_{B_{\hat{r}}} \mathcal{B}_h^n \left(\nabla \Delta_h^n u, \nabla (\eta^2 \Delta_h^n u [\Gamma_n^{(h)} - k]^+) \right) dx \\ &= \int_{B_{\hat{r}}} \eta^2 [\Gamma_n^{(h)} - k]^+ \mathcal{B}_h^n (\nabla \Delta_h^n u, \nabla \Delta_h^n u) dx + \int_{B_{\hat{r}} \cap [\Gamma_n^{(h)} \geq k]} \eta^2 \mathcal{B}_h^n (\nabla \Delta_h^n u, \nabla \Gamma_n^{(h)}) \Delta_h^n u dx \\ &\quad + \int_{B_{\hat{r}}} 2\eta \mathcal{B}_h^n (\nabla \Delta_h^n u, \nabla \eta) [\Gamma_n^{(h)} - k]^+ \Delta_h^n u dx \\ &=: T_1 + T_2 + T_3.\end{aligned}$$

T_1 is non-negative, and clearly

$$\begin{aligned}T_2 &= \frac{1}{2} \int_{B_{\hat{r}} \cap [\Gamma_n^{(h)} \geq k]} \eta^2 \mathcal{B}_h^n (\nabla \Gamma_n^{(h)}, \nabla \Gamma_n^{(h)}) dx, \\ T_3 &= \int_{B_{\hat{r}} \cap [\Gamma_n^{(h)} \geq k]} \eta \mathcal{B}_h^n (\nabla \Gamma_n^{(h)}, \nabla \eta) (\Gamma_n^{(h)} - k) dx,\end{aligned}$$

hence

$$\int_{B_{\hat{r}} \cap [\Gamma_n^{(h)} \geq k]} \eta^2 \mathcal{B}_h^n (\nabla \Gamma_n^{(h)}, \nabla \Gamma_n^{(h)}) dx \leq -2 \int_{B_{\hat{r}} \cap [\Gamma_n^{(h)} \geq k]} \eta \mathcal{B}_h^n (\nabla \Gamma_n^{(h)}, \nabla \eta) (\Gamma_n^{(h)} - k) dx.$$

The same arguments leading from (3.4) to (3.5) then show

$$\int_{B_{\hat{r}} \cap [\Gamma_n^{(h)} \geq k]} \eta^2 \mathcal{B}_h^n (\nabla \Gamma_n^{(h)}, \nabla \Gamma_n^{(h)}) dx \leq c \int_{B_{\hat{r}} \cap [\Gamma_n^{(h)} \geq k]} \mathcal{B}_h^n (\nabla \eta, \nabla \eta) (\Gamma_n^{(h)} - k)^2 dx. \quad (3.8)$$

Again by (3.1) it is immediate that

$$\text{r.h.s. of (3.8)} \xrightarrow{h \rightarrow 0} c \int_{B_{\hat{r}} \cap [\Gamma_n \geq k]} D^2 F(\nabla u) (\nabla \eta, \nabla \eta) (\Gamma_n - k)^2 dx,$$

whereas on the l.h.s. of (3.8) we observe that the integrand is ≥ 0 and pointwise convergent (since $u \in W_{2,loc}^2(\Omega)$), thus Fatou's lemma is applicable, and we deduce from (3.8)

$$\int_{B_{\hat{r}} \cap [\Gamma_n \geq k]} \eta^2 D^2 F(\nabla u) (\nabla \Gamma_n, \nabla \Gamma_n) dx \leq c \int_{B_{\hat{r}} \cap [\Gamma_n \geq k]} D^2 F(\nabla u) (\nabla \eta, \nabla \eta) (\Gamma_n - k)^2 dx. \quad (3.9)$$

Recalling the ellipticity estimates for f and g we get from (3.9) the Caccioppoli-type inequality

$$\begin{aligned}\int_{B_{\hat{r}} \cap [\Gamma_n \geq k]} \eta^2 |\nabla \Gamma_n|^2 dx &\leq c \left[\int_{B_{\hat{r}} \cap [\Gamma_n \geq k]} \tilde{\Gamma}^{\frac{s-2}{2}} (\Gamma_n - k)^2 |\nabla \eta|^2 dx \right. \\ &\quad \left. + \int_{B_{\hat{r}} \cap [\Gamma_n \geq k]} \Gamma_n^{\frac{t-2}{2}} (\Gamma_n - k)^2 |\nabla \eta|^2 dx \right].\end{aligned} \quad (3.10)$$

Let $\eta = 1$ on B_r , $|\nabla\eta| \leq c/(\hat{r} - r)$, $0 \leq \eta \leq 1$. Let us further abbreviate

$$A(k, r) := B_r \cap [\Gamma_n \geq k].$$

We proceed similar to [Bi], proof of Theorem 5.22: we have

$$\begin{aligned} \int_{A(k,r)} (\Gamma_n - k)^{\frac{n}{n-1}} dx &\leq \int_{B_{\hat{r}}} \left[\eta(\Gamma_n - k)^+ \right]^{\frac{n}{n-1}} dx \\ &\leq c \left[\int_{B_{\hat{r}}} |\nabla(\eta(\Gamma_n - k)^+)| dx \right]^{\frac{n}{n-1}} \\ &\leq c [I_1^{\frac{n}{n-1}} + I_2^{\frac{n}{n-1}}], \end{aligned} \quad (3.11)$$

$$\begin{aligned} I_1^{\frac{n}{n-1}} &:= \left[\int_{A(k,r)} |\nabla\eta|(\Gamma_n - k) dx \right]^{\frac{n}{n-1}} \\ &\leq c(\hat{r} - r)^{-\frac{n}{n-1}} \left[\int_{A(k,\hat{r})} \Gamma_n^{\frac{t-2}{4}} (\Gamma_n - k) \Gamma_n^{\frac{2-t}{4}} dx \right]^{\frac{n}{n-1}} \\ &\leq c(\hat{r} - r)^{-\frac{n}{n-1}} \left[\int_{A(k,\hat{r})} \Gamma_n^{\frac{t-2}{2}} (\Gamma_n - k)^2 dx \right]^{\frac{1}{2} \frac{n}{n-1}} \left[\int_{A(k,\hat{r})} \Gamma_n^{\frac{2-t}{2}} dx \right]^{\frac{1}{2} \frac{n}{n-1}}, \end{aligned} \quad (3.12)$$

$$\begin{aligned} I_2^{\frac{n}{n-1}} &:= \left[\int_{A(k,\hat{r})} \eta |\nabla\Gamma_n| dx \right]^{\frac{n}{n-1}} \\ &\leq c \left[\int_{A(k,\hat{r})} \eta^2 |\nabla\Gamma_n|^2 dx \right]^{\frac{1}{2} \frac{n}{n-1}} |A(k, \hat{r})|^{\frac{1}{2} \frac{n}{n-1}}, \end{aligned} \quad (3.13)$$

where in (3.11) we made use of Sobolev's inequality, whereas (3.12) and (3.13) just follow from Hölder's inequality. Inserting (3.10) into the r.h.s. of (3.13), we get

$$\begin{aligned} &\int_{A(k,r)} (\Gamma_n - k)^{\frac{n}{n-1}} dx \\ &\leq c(\hat{r} - r)^{-\frac{n}{n-1}} \left\{ \left[\int_{A(k,\hat{r})} \Gamma_n^{\frac{t-2}{2}} (\Gamma_n - k)^2 dx \right]^{\frac{1}{2} \frac{n}{n-1}} \left[\int_{A(k,\hat{r})} \Gamma_n^{\frac{2-t}{2}} dx \right]^{\frac{1}{2} \frac{n}{n-1}} \right. \\ &\quad \left. + \left[\int_{A(k,\hat{r})} \Gamma_n^{\frac{t-2}{2}} (\Gamma_n - k)^2 dx + \int_{A(k,\hat{r})} \tilde{\Gamma}^{\frac{s-2}{2}} (\Gamma_n - k)^2 dx \right]^{\frac{1}{2} \frac{n}{n-1}} |A(k, \hat{r})|^{\frac{1}{2} \frac{n}{n-1}} \right\}. \end{aligned} \quad (3.14)$$

Let $\nu, \varkappa > 1$ be arbitrary for the moment. We recall (3.1) and apply Hölder's inequality to obtain

$$\begin{aligned} \int_{A(k,r)} \Gamma_n^{\frac{t-2}{2}} (\Gamma_n - k)^2 dx &\leq \int_{A(k,r)} (\Gamma_n - k)^{\frac{n}{n-1} \frac{1}{\nu}} \Gamma_n^{2 - \frac{n}{n-1} \frac{1}{\nu}} \Gamma_n^{\frac{t-2}{2}} dx \\ &\leq c_1(\nu) \left[\int_{A(k,r)} (\Gamma_n - k)^{\frac{n}{n-1}} dx \right]^{\frac{1}{\nu}}, \end{aligned} \quad (3.15)$$

$$\begin{aligned}
\int_{A(k,r)} \tilde{\Gamma}^{\frac{s-2}{2}} (\Gamma_n - k)^2 dx &= \int_{A(k,r)} (\Gamma_n - k)^{\frac{n}{n-1} \frac{1}{\nu}} (\Gamma_n - k)^{2 - \frac{n}{n-1} \frac{1}{\nu}} \tilde{\Gamma}^{\frac{s-2}{2}} dx \\
&\leq \int_{A(k,r)} (\Gamma_n - k)^{\frac{n}{n-1} \frac{1}{\nu}} \Gamma_n^{2 - \frac{n}{n-1} \frac{1}{\nu}} \tilde{\Gamma}^{\frac{s-2}{2}} dx \\
&\leq c_1(\nu) \left[\int_{A(k,r)} (\Gamma_n - k)^{\frac{n}{n-1}} dx \right]^{\frac{1}{\nu}}, \tag{3.16}
\end{aligned}$$

$c_1(\nu)$ denoting a local constant depending on our fixed ball $B_R(x_0)$. In the same way it follows

$$\int_{A(k,\hat{r})} \Gamma_n^{\frac{2-t}{2}} dx = \int_{A(k,\hat{r})} \Gamma_n^{\frac{1}{\varkappa} \frac{t-2}{2}} \Gamma_n^{2 - \frac{1}{\varkappa} \frac{t-2}{2}} dx \leq c_2(\varkappa) \left[\int_{A(k,\hat{r})} \Gamma_n^{\frac{t-2}{2}} dx \right]^{\frac{1}{\varkappa}}, \tag{3.17}$$

$$|A(k,\hat{r})| \leq \int_{A(k,\hat{r})} \Gamma_n^{\frac{1}{\varkappa} \frac{t-2}{2}} dx \leq c_2(\varkappa) \left[\int_{A(k,\hat{r})} \Gamma_n^{\frac{t-2}{2}} dx \right]^{\frac{1}{\varkappa}}. \tag{3.18}$$

This gives

$$\begin{aligned}
\tau(k,r) &:= \int_{A(k,r)} \left(\Gamma_n^{\frac{t-2}{2}} (\Gamma_n - k)^2 + \tilde{\Gamma}^{\frac{s-2}{2}} (\Gamma_n - k)^2 \right) dx \\
&\leq c_1(\nu) \left[\int_{A(k,r)} (\Gamma_n - k)^{\frac{n}{n-1}} dx \right]^{\frac{1}{\nu}} \\
&\leq c_3(\nu, \varkappa) (\hat{r} - r)^{-\frac{1}{\nu} \frac{n}{n-1}} \tau(k, \hat{r})^{\frac{1}{\nu} \frac{1}{2} \frac{n}{n-1}} \left[\int_{A(k,\hat{r})} \Gamma_n^{\frac{t-2}{2}} dx \right]^{\frac{1}{\varkappa \nu} \frac{1}{2} \frac{n}{n-1}},
\end{aligned}$$

where we made use of (3.15), (3.16) to obtain the first inequality, (3.14), (3.17) and (3.18) imply the second one. If we abbreviate

$$a(k,r) := \int_{A(k,r)} \Gamma_n^{\frac{t-2}{2}} dx,$$

then it is shown that for any $k > 0$ and for all $\nu, \varkappa > 1$, $0 < r < \hat{r} \leq R$ the inequality

$$\tau(k,r) \leq c_3 (\hat{r} - r)^{-\frac{1}{\nu} \frac{n}{n-1}} \tau(k, \hat{r})^{\frac{1}{\nu} \frac{1}{2} \frac{n}{n-1}} a(k, \hat{r})^{\frac{1}{\varkappa \nu} \frac{1}{2} \frac{n}{n-1}} \tag{3.19}$$

is valid. Obviously $a(h, \hat{r}) \leq (h - k)^{-2} \tau(k, \hat{r})$, if $h > k$, and from (3.19) (replacing k by $h > k$) we get

$$\begin{aligned}
\tau(h,r) &\leq c_3 (\hat{r} - r)^{-\gamma} \tau(h, \hat{r})^{\frac{1}{\nu} \frac{1}{2} \frac{n}{n-1}} a(h, \hat{r})^{\frac{1}{\varkappa \nu} \frac{1}{2} \frac{n}{n-1}} \\
&\leq c_3 (\hat{r} - r)^{-\gamma} (h - k)^{-\beta} \tau(h, \hat{r})^{\frac{1}{\nu} \frac{1}{2} \frac{n}{n-1}} \tau(k, \hat{r})^{\frac{1}{\varkappa \nu} \frac{1}{2} \frac{n}{n-1}}
\end{aligned}$$

with suitable positive exponents γ, β . Since $\tau(h, \hat{r}) \leq \tau(k, \hat{r})$, we finally arrive at

$$\tau(h,r) \leq c_3 (\hat{r} - r)^{-\gamma} (h - k)^{-\beta} \tau(k, \hat{r})^{\frac{1}{2} \frac{n}{n-1} \frac{1}{\nu} [1 + \frac{1}{\varkappa}]}, \quad 0 < k < h, \quad 0 < r < \hat{r} \leq R. \tag{3.20}$$

In (3.20) ν and \varkappa are still in our disposal, and for $\nu, \varkappa > 1$ but very close to 1 we can arrange that

$$\frac{1}{2} \frac{n}{n-1} \frac{1}{\nu} \left[1 + \frac{1}{\varkappa} \right] > 1,$$

thus a lemma of Stampacchia [St, Lemma 5.1, p.219] can be applied to (3.20) implying $\Gamma_n \in L^\infty(B_{R/2}(x_0))$. Since the ball $B_R(x_0)$ was arbitrary, it follows that $\partial_n u$ is in the space $L^\infty_{loc}(\Omega)$.

Next we fix a coordinate direction e_m , $m < n$, and let $\Gamma_m := 1 + |\partial_m u|^2$, $A(k, r) := B_r \cap [\Gamma_m \geq k]$ etc. We indicate the changes in the foregoing calculations: (3.9) holds for Γ_m in place of Γ_n , and if we use $\partial_n u \in L^\infty_{loc}(\Omega)$, then (3.10) can be replaced by

$$\int_{A(k, \hat{r})} \eta^2 |\nabla \Gamma_m|^2 dx \leq c \int_{A(k, \hat{r})} \tilde{\Gamma}^{\frac{s-2}{2}} (\Gamma_m - k)^2 |\nabla \eta|^2 dx. \quad (3.10')$$

Here we just observed $\Gamma_n^{\frac{t-2}{2}} \leq c \tilde{\Gamma}^{\frac{s-2}{2}}$. (3.11) remains valid for Γ_m , and we have

$$I_1^{\frac{n}{n-1}} \leq c(\hat{r} - r)^{-\frac{n}{n-1}} \left[\int_{A(k, \hat{r})} \Gamma_m^{\frac{s-2}{2}} (\Gamma_m - k)^2 dx \right]^{\frac{1}{2} \frac{n}{n-1}} \left[\int_{A(k, \hat{r})} \Gamma_m^{\frac{2-s}{2}} dx \right]^{\frac{1}{2} \frac{n}{n-1}}. \quad (3.12')$$

(3.13) of course is true for Γ_m , and as before we can use (3.10') on the r.h.s. of the Γ_m -version of (3.13), which leads in combination with (3.12') and (3.13) to

$$\begin{aligned} & \int_{A(k, r)} (\Gamma_m - k)^{\frac{n}{n-1}} dx \\ & \leq c(\hat{r} - r)^{-\frac{n}{n-1}} \left\{ \left[\int_{A(k, \hat{r})} \Gamma_m^{\frac{s-2}{2}} (\Gamma_m - k)^2 dx \right]^{\frac{1}{2} \frac{n}{n-1}} \left[\int_{A(k, \hat{r})} \Gamma_m^{\frac{2-s}{2}} dx \right]^{\frac{1}{2} \frac{n}{n-1}} \right. \\ & \quad \left. + \left[\int_{A(k, \hat{r})} \tilde{\Gamma}^{\frac{s-2}{2}} (\Gamma_m - k)^2 dx \right]^{\frac{1}{2} \frac{n}{n-1}} |A(k, \hat{r})|^{\frac{1}{2} \frac{n}{n-1}} \right\}. \end{aligned} \quad (3.14')$$

Obviously (see (3.15), (3.16)) it holds for $\nu > 1$ (recall (3.1)!)

$$\int_{A(k, r)} \tilde{\Gamma}^{\frac{s-2}{2}} (\Gamma_m - k)^2 dx \leq c_1(\nu) \left[\int_{A(k, r)} (\Gamma_m - k)^{\frac{n}{n-1}} dx \right]^{\frac{1}{\nu}}, \quad (3.16')$$

and in accordance with (3.17), (3.18) we find

$$\int_{A(k, \hat{r})} \Gamma_m^{\frac{2-s}{2}} dx \leq c_2(\varkappa) \left[\int_{A(k, \hat{r})} \Gamma_m^{\frac{s-2}{2}} dx \right]^{\frac{1}{\varkappa}}, \quad (3.17')$$

$$|A(k, \hat{r})| \leq c_2(\varkappa) \left[\int_{A(k, \hat{r})} \Gamma_m^{\frac{s-2}{2}} dx \right]^{\frac{1}{\varkappa}}. \quad (3.18')$$

This gives by (3.14'), (3.16') and (3.17'), (3.18'), respectively,

$$\begin{aligned} & \int_{A(k, r)} \tilde{\Gamma}^{\frac{s-2}{2}} (\Gamma_m - k)^2 dx \\ & \leq c_1(\nu) (\hat{r} - r)^{-\frac{1}{\nu} \frac{n}{n-1}} \left[\int_{A(k, \hat{r})} \tilde{\Gamma}^{\frac{s-2}{2}} (\Gamma_m - k)^2 dx \right]^{\frac{1}{2} \frac{1}{\nu} \frac{n}{n-1}} \\ & \quad \cdot \left\{ \left[\int_{A(k, \hat{r})} \Gamma_m^{\frac{2-s}{2}} dx \right]^{\frac{1}{2} \frac{n}{n-1}} + |A(k, \hat{r})|^{\frac{1}{2} \frac{n}{n-1}} \right\}^{\frac{1}{\nu}} \\ & \leq c_3(\nu, \varkappa) (\hat{r} - r)^{-\frac{1}{\nu} \frac{n}{n-1}} \left[\int_{A(k, \hat{r})} \tilde{\Gamma}^{\frac{s-2}{2}} (\Gamma_m - k)^2 dx \right]^{\frac{1}{2} \frac{1}{\nu} \frac{n}{n-1}} \left[\int_{A(k, \hat{r})} \Gamma_m^{\frac{s-2}{2}} dx \right]^{\frac{1}{2} \frac{1}{\varkappa} \frac{1}{\nu} \frac{n}{n-1}}. \end{aligned}$$

Let us now define

$$\begin{aligned}\tau'(k, r) &:= \int_{A(k, r)} \tilde{\Gamma}^{\frac{s-2}{2}} (\Gamma_m - k)^2 dx, \\ a'(k, r) &:= \int_{A(k, r)} \Gamma_m^{\frac{s-2}{2}} dx.\end{aligned}$$

Then the foregoing inequality implies (3.19) for the quantities τ' , a' , and since again ($h > k$)

$$a'(h, \hat{r}) \leq (h - k)^{-2} \tau'(k, \hat{r})$$

we deduce (3.20) for τ' which means $\Gamma_m \in L^\infty(B_{R/2}(x_0))$, thus $\partial_m u \in L^\infty_{loc}(\Omega)$. \square

Remark 3.1. *The information $\partial_n u \in L^\infty_{loc}(\Omega)$ is not really needed for the proof of $\tilde{\nabla} u \in L^\infty_{loc}(\Omega, \mathbb{R}^{n-1})$. If we keep the quantity*

$$\int_{A(k, \hat{r})} \Gamma_n^{\frac{t-2}{2}} (\Gamma_m - k)^2 dx$$

on the r.h.s. of (3.10'), then this item occurs on the r.h.s. of (3.14') (it has to be added to $\int_{A(k, \hat{r})} \tilde{\Gamma}^{\frac{s-2}{2}} (\Gamma_m - k)^2 dx$). On the other hand we have (compare (3.16'))

$$\int_{A(k, r)} \Gamma_n^{\frac{t-2}{2}} (\Gamma_m - k)^2 dx \leq c_1(\nu) \left[\int_{A(k, r)} (\Gamma_m - k)^{\frac{n}{n-1}} dx \right]^{1/\nu},$$

hence

$$\begin{aligned}\tau''(k, r) &:= \int_{A(k, r)} \left[\tilde{\Gamma}^{\frac{s-2}{2}} + \Gamma_n^{\frac{t-2}{2}} \right] (\Gamma_m - k)^2 dx \\ &\leq c_1(\nu) \left[\int_{A(k, r)} (\Gamma_m - k)^{\frac{n}{n-1}} dx \right]^{1/\nu} \\ &\leq c_1(\nu) (\hat{r} - r)^{-\frac{1}{\nu} \frac{n}{n-1}} \tau''(k, \hat{r})^{\frac{1}{2} \frac{1}{\nu} \frac{n}{n-1}} \left\{ \left[\int_{A(k, \hat{r})} \Gamma_m^{\frac{2-s}{2}} dx \right]^{\frac{1}{2} \frac{n}{n-1}} + |A(k, \hat{r})|^{\frac{1}{2} \frac{n}{n-1}} \right\}^{\frac{1}{\nu}}\end{aligned}$$

and the terms in $\{\dots\}$ are treated via (3.17'), (3.18'). This gives (3.19) for τ'' and a' , and from $a'(h, \hat{r}) \leq (h - k)^{-2} \tau''(k, \hat{r})$, $h > k$, the τ'' -version of (3.20) follows which again gives $\partial_m u \in L^\infty_{loc}(\Omega)$.

References

- [Ad] Adams, R. A., Sobolev spaces. Academic Press, New York-San Francisco-London 1975.
- [Bi] Bildhauer, M., Convex variational problems: linear, nearly linear and anisotropic growth conditions. Lecture Notes in Mathematics 1818, Springer, Berlin-Heidelberg-New York, 2003.
- [BF1] Bildhauer, M., Fuchs, M., Elliptic variational problems with nonstandard growth. International Mathematical Series, Vol. 1, Nonlinear problems in mathematical physics and related topics I, in honor of Prof. O.A. Ladyzhenskaya. By Tamara Rozhkovskaya, Novosibirsk, Russia, March 2002 (in Russian), 49–62. By Kluwer/Plenum Publishers, June 2002 (in English), 53–66.

- [BF2] Bildhauer, M., Fuchs, M., Higher integrability of the gradient for vectorial minimizers of decomposable variational integrals. Submitted.
- [BFM] Bildhauer, M., Fuchs, M., Mingione, G., A priori gradient bounds and local $C^{1,\alpha}$ -estimates for (double) obstacle problems under nonstandard growth conditions. *Z. Anal. Anw.* 20, 959–985 (2001).
- [BFZ] Bildhauer, M., Fuchs, M., Zhong, X., Variational integrals with a wide range of anisotropy. To appear in *Algebra i Analiz*.
- [Ch] Choe, H. J., Interior behaviour of minimizers for certain functionals with nonstandard growth. *Nonlinear Analysis, Theory, Methods & Appl.* 19.10 (1992), 933-945.
- [ELM] Esposito, L., Leonetti, F., Mingione, G., Regularity for minimizers of functionals with p-q growth. *Nonlinear Diff. Equ. Appl.* 6 (1999), 133-148.
- [FS] Fusco, N., Sbordone, C., Some remarks on the regularity of minima of anisotropic integrals. *Comm. P.D.E.* 18, 153–167 (1993).
- [Gi] Giaquinta, M., Growth conditions and regularity, a counterexample. *Manus. Math.* 59 (1987), 245–248.
- [GT] Gilbarg, D., Trudinger, N.S., Elliptic partial differential equations of second order. *Grundlehren der math. Wiss.* 224, second ed., revised third print., Springer, Berlin-Heidelberg-New York 1998.
- [Ho] Hong, M.C., Some remarks on the minimizers of variational integrals with non standard growth conditions. *Boll. U.M.I.* (7) 6-A (1992), 91–101.
- [Ma1] Marcellini, P., Regularity of minimizers of integrals of the calculus of variations with non standard growth conditions. *Arch. Rat. Mech. Anal.* 105 (1989), 267–284.
- [Ma2] Marcellini, P., Everywhere regularity for a class of elliptic systems without growth conditions. *Ann. Scuola Norm. Sup. Pisa* 23 (1996), 1–25.
- [MM] Massari, U., Miranda, M., Minimal surfaces of codimension one. *North-Holland Mathematics Studies* 91, North-Holland, Amsterdam-New York-Oxford, 1983.
- [St] Stampacchia, G., Le problème de Dirichlet pour les équations elliptiques du second ordre á coefficients discontinus. *Ann. Inst. Fourier Grenoble* 15.1 (1965), 189–258.
- [UU] Ural'tseva, N.N., Urdaletova, A.B., The boundedness of the gradients of generalized solutions of degenerate quasilinear nonuniformly elliptic equations. *Vestn. Leningr. Univ.* 1983, Mat. Mekh. Astron. no. 4 (1983), 50–56 (in Russian). English translation: *Vestn. Leningr. Univ. Math* 16 (1984), 263–270.